

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 April 2003 (17.04.2003)

PCT

(10) International Publication Number
WO 03/031632 A1

(51) International Patent Classification⁷: **C12N 15/82**

(21) International Application Number: **PCT/EP02/11188**

(22) International Filing Date: **2 October 2002 (02.10.2002)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
01203760.2 **5 October 2001 (05.10.2001)** **EP**

(71) Applicant (*for all designated States except US*):
**VLAAMS INTERUNIVERSITAIR INSTITUUT
VOOR BIOTECHNOLOGIE VZW** [BE/BE]; Rijviss-
chestraat 120, B-9052 Zwijnaarde (BE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **DEPICKER, Anna**
[BE/BE]; Molenstraat 61, B-9820 Merelbeke (BE). **VAN
HOUDT, Helena** [BE/BE]; Lieskensweg 7, B-9080
Lochristi (BE).

(74) Common Representative: **VLAAMS INTERUNIVER-
SITAIR INSTITUUT VOOR BIOTECHNOLOGIE
VZW**; Rijvisschestraat 120, B-9052 Zwijnaarde (BE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/031632 A1

(54) Title: **AN EFFICIENT SYSTEM FOR RNA SILENCING**

(57) Abstract: The invention relates to a method for efficient RNA silencing of target genes in eucaryotic cells, particularly plant cells. Consequently, the method can be used to reduce the phenotypic expression of an endogenous gene in a plant cell. Furthermore the method can be applied in a high throughput screening for mutant phenotypes as a result of RNA silencing of any endogene.

An efficient system for RNA silencing

Field of the invention

The invention relates to a method for efficient RNA silencing in eucaryotic cells, particularly plant cells. Consequently, the method can be used to reduce the phenotypic expression of an endogenous gene in a plant cell. Furthermore the method can be applied in a high throughput screening for RNA silencing.

Background of the invention

RNA silencing is a type of gene regulation based on sequence-specific targeting and degradation of RNA. The term encompasses related pathways found in a broad range of eukaryotic organisms, including fungi, plants, and animals. In plants, RNA silencing serves as an antiviral defense, and many plant viruses encode suppressors of silencing. Also it becomes clear that elements of the RNA silencing system are essential for gene regulation in development. The emerging view is that RNA silencing is part of a sophisticated network of interconnected pathways for cellular defense, transposon surveillance, and regulation of development. Based on the sequence specific RNA degradation, RNA silencing has become a powerful tool to manipulate gene expression experimentally. RNA silencing was first discovered in transgenic plants, where it was termed co-suppression or posttranscriptional gene silencing (PTGS). Sequence-specific RNA degradation processes related to PTGS have also been found in ciliates, fungi, and a variety of animals from *Caenorhabditis elegans* to mice (RNA interference). A key feature uniting the RNA silencing pathways in different organisms is the importance of double-stranded RNA (dsRNA) as a trigger or an intermediate. The dsRNA is cleaved into small interfering RNAs (21 to 25 nucleotides) of both polarities, and these are thought to act as guides to direct the RNA degradation machinery to the target RNAs. An intriguing aspect of RNA silencing in plants is that it can be triggered locally and then spread via a mobile silencing signal. In plants, RNA silencing is correlated with methylation of homologous transgene DNA in the nucleus. Other types of epigenetic modifications may be associated with silencing in other organisms.

It is known from the art that transgenes encoding ds or self-complementary (hairpin) RNAs of endogenous gene sequences are highly effective at directing the cell's degradation mechanism against endogenous (ss) mRNAs, thus giving targeted gene

suppression. This discovery has enabled the transgenic enhancement of a plant's defense mechanism against viruses that it is unable to combat unaided. It has also shed light on how antisense and co-suppression might operate: by the inadvertent integration of two copies of the transgenes in an inverted repeat orientation, such that
5 read-through transcription from one gene into the adjacent copy produces RNA with self-complementary sequences.

RNA silencing is induced in plants by transgenes designed to produce either sense or antisense transcripts. Furthermore, transgenes engineered to produce self-complementary transcripts (dsRNAs) are potent and consistent inducers of RNA
10 silencing. Finally, replication of plant viruses, many of which produce dsRNA replication intermediates, causes a type of RNA silencing called Virus Induced Gene Silencing (VIGS). Whether VIGS, and the different types of transgene-induced RNA silencing in plants result from similar or distinct mechanisms is still a matter of debate. However, recent genetic evidence raises the possibility that the RNA silencing pathway
15 is branched and that the branches converge in the production of dsRNA.

Until recently RNA silencing was viewed primarily as a thorn in the side of plant molecular geneticists, limiting expression of transgenes and interfering with a number of applications that require consistent, high-level transgene expression. With our present understanding of the process, however, it is clear that RNA silencing could
20 have enormous potential for engineering control of gene expression, as well as for the use as a tool in functional genomics. It could be experimentally induced and targeted to a single specific gene or even to a family of related genes. Likewise, ds RNA-induced TGS may have similar potential to control gene expression. Although several methods for RNA silencing have been described in the art (WO99/53050,
25 WO99/32619, WO99/61632, and W098/53083), there is clearly a need to develop alternative and more efficient tools for RNA silencing. In the present invention we have developed a highly efficient method for RNA silencing that can also be used as a tool for high throughput silencing. Said method uses a host that carries already a silenced locus and a second recombinant gene comprising a region that is homologous with the
30 silenced locus. Although it is known from the art that the recombinant gene will be silenced, we have surprisingly found that also target genes, which have no significant homology with the silenced locus but have homology with the recombinant gene, are efficiently silenced.

Figure legends

Fig. 1: Schematical outline of homology between a silenced locus X, a recombinant gene Y and a target gene Z.

- 5 Fig. 2: Schematical outline of the T-DNA constructs that are present in silenced locus X₁, recombinant gene Y₁ and target gene Z₁ (T-DNAs of pGVCHS287, pGUSchsS and pXD610 respectively) and of the transcript homology between X₁, Y₁ and Z₁.

- 10 LB and RB: left and right T-DNA border respectively; Pnos: nopaline synthase promoter; hpt: hygromycin phosphotransferase coding sequence; 3'nos: 3'untranslated region of the nopaline synthase gene; P35S: Cauliflower mosaic virus 35S promoter; nptII c.s., neomycin phosphotransferase II coding sequence; 3'chs: 3'untranslated region of the chalcone synthase gene of *Anthirrium majus*; +1: transcription start; A_n: poly A-tail; gus c.s.: β-glucuronidase coding sequence; Pss: promoter of the small
15 subunit of rubisco; bar: phosphinotricine transferase coding sequence; 3'g7: 3'untranslated region of the *Agrobacterium octopine* T-DNA gene 7; 3'ocs: 3'untranslated region of octopine synthase gene.

- Fig. 3: Schematical outline of the T-DNA construct present in silenced locus X₁ and of
20 the transiently introduced T-DNAs Y₂ (T-DNAs of pGVCHS287 and pPs35SCAT1S3chs, respectively) and of the transcript homology between X₁, Y₂ and Z₂ (the catalase1 endogene). Abbreviations as in Fig. 2

- Fig. 4: Schematical outline of the T-DNA constructs present in silenced locus X₂ and of
25 the transiently introduced T-DNAs Y₂ (T-DNAs of pGUSchsS + pGUSchsAS, and pPs35SCAT1S3chs, respectively) and of the transcript homology between X₂, Y₂ and Z₂ (the catalase1 endogene). Abbreviations as in Fig. 2

Fig.5: pPs35SCAT1S3chs

30

Detailed description of the invention

The present invention deals with an efficient method for RNA silencing in an eucaryotic host. The method makes use of a host that already comprises a silenced locus. Such a silenced locus can for example be generated by methods known in the art. For

example the publication of De Buck and Depicker, 2001 and other publications, and also patents WO99/53050, WO99/32619, WO99/61632, and W098/53083 describe methods to obtain RNA silencing and for generating a silenced recombinant locus. The 'target gene' is here defined as the gene of interest for silencing or to down-regulate its expression. An important aspect of this invention is that said target gene has no significant homology with the silenced locus. No significant homology means that either the overall homology is less than 40, 35, 30, 25% or even less, or that no contiguous stretch of at least 23 identical nucleotides are present (Thomas et al., 2001). Homology is typically measured using sequence analysis software (e.g., Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, Wis. 53705). Such software matches similar sequences by assigning degrees of homology to various insertions, deletions, substitutions, and other modifications. Silencing of said target gene in the present invention occurs via an intermediate step and hence our method is designated as domino silencing (Fig. 1). In said intermediate step a recombinant gene construct is introduced by transformation into the host comprising the silenced locus. Said recombinant gene construct has a region of homology with the silenced locus already present. Said region of homology is preferably more than 60, 70, 80, 90, 95 or even more than 99% homologous. The homologous region between the silenced locus and said recombinant gene can be found in the 5' untranslated or 3' untranslated region of the recombinant gene construct. Furthermore, said recombinant gene construct has a region of minimal 23 nucleotides (Thomas et al., 2001), but preferably longer, that are identical with the target gene, or has a region of overall homology of more than 60, 70, 80, 90, 95 or even more than 99%. A recombinant gene is defined herein as a construct which does not naturally occur in nature. A non-limiting example of a recombinant gene construct is a construct wherein the coding region of a gene is operably linked to a 5' untranslated region and/or to a 3' untranslated region of one or more other genes, alternatively said 5' or 3' untranslated region is an artificial sequence.

Thus in one embodiment the invention provides a method for obtaining efficient RNA silencing of a target gene comprising the introduction of a recombinant gene into a host that comprises a silenced locus and an unsilenced target gene whereby said recombinant gene comprises a region that is homologous with said silenced locus and

whereby said target gene has homology with said recombinant gene but has no significant homology with said silenced locus.

In another embodiment the method is used wherein said host is a plant or plant cell.

In another embodiment the method of the invention can be used for high throughput gene silencing. Indeed, a recombinant gene library can be made wherein for example every gene or coding region thereof is combined with (operably linked with) a region of homology with the silenced gene that resides in the silenced locus and said recombinant gene library can be transformed to an eukaryotic host or individual (specific) genes derived from said recombinant gene library can be transformed into an eukaryotic host wherein silencing of specific genes is wanted.

In yet another embodiment the invention provides a plant or plant cell that comprises a silenced locus and wherein a silenced target gene is obtained through the introduction of a recombinant gene according to the current method of the invention.

In yet another embodiment the RNA silencing of the target gene is obtained in more than 80, 85, 90 or 95% of the transgenic organisms.

In yet another embodiment the RNA silencing of the target gene occurs at an efficiency of more than 80, 85, 90 or 95 % as compared to the level of the unsilenced expression of the target gene.

20

Examples

A posttranscriptionally silenced inverted repeat transgene locus can trigger silencing of a reporter gene producing non-homologous transcripts.

We studied the interaction between three transgene loci X_1 , Y_1 and Z_1 (Fig. 2, For a detailed description of all loci and constructs, see materials and methods) to address the question whether or not a stepwise homology between loci can lead to silencing.

It has been demonstrated previously that the posttranscriptionally silenced *nptII* genes in locus X_1 are capable to in trans silence transiently expressed genes with partial transcript homology to their *nptII* transcripts (Van Houdt et al., 2000 b). We subsequently found that also a stably expressed β -glucuronidase (*gus*) gene (in locus Y_1), with partial transcript homology to the *nptII* transcripts of the silencing inducing locus X_1 , becomes efficiently silenced in trans (Fig. 2: X_1 and Y_1 and table 1: X_1Y_1 compared to Y_1). On the contrary, the *nptII* genes of locus X_1 are not able to trigger silencing of the *gus* genes in locus Z_1 which is expected as the genes of both loci produce transcripts without significant homology (Fig. 2). The homology between the

two transcripts of X_1 and Y_1 is mainly situated in the 3'untranslated region (250 nucleotides), but also the 5'untranslated sequences show a small region of homology (29 nucleotides). These results demonstrate that the in trans silencing effects are not triggered by promoter homology. When Y_1 and Z_1 loci are combined in so called Y_1Z_1 hybrids both types of gus genes, having transcript homology in the gus coding sequence of 1809 nucleotides, remain highly expressed as reflected in the normal gus activity showing that the RNA silencing mechanism does not become activated (Table 1: Y_1Z_1 compared to Y_1 and Z_1). Surprisingly, upon creation of a stepwise homology between X_1 and Z_1 by introducing locus Y_1 , the new observation described here is that also the gus expression in locus Z_1 is reduced in $X_1Y_1Z_1$ plants (Table 1: $X_1Y_1Z_1$ compared to Y_1Z_1). Thus, creating a stepwise homology between a silenced locus and a target gene by introducing a recombinant gene is sufficient to trigger silencing of the target.

15 Silencing inducing transgene loci can trigger silencing of a non-homologous endogene.

We further assessed the universality and the usefulness in high throughput functional gene analyses of silencing elicited by a stepwise homology in trans, called domino silencing. Therefore, we evaluated whether the expression of the tobacco endogenous catalase1 (cat1) genes is reduced in plants carrying a silencing locus (X locus) showing no significant homology with the catalase endogene by introducing a recombinant gene (Y construct). As silencing locus we used either X_1 or X_2 (Fig.2: locus X_1 , Fig.3: locus X_2), in either case containing the 3' chalcone synthase sequences of *Anthirrinum majus* (3'chs). As transmitter for silencing we constructed a recombinant gene composed of the catalase1 coding sequence and the 3' chs region under control of the 35S promoter (P35S) (residing on T-DNA pPs35SCAT1S3chs, Fig.2 and 3: T-DNA in Y_2). The recombinant cat1 3'chs genes (Y_2) were introduced in tobacco leaves bearing locus X_1 (or X_2) via *Agrobacterium* injection. As a negative control, we introduced a recombinant gene in which the cat1 coding sequence is replaced by the gus coding sequence (pGUSchsS, T-DNA construct as in locus Y_1 Fig.1). In this case, no stepwise homology is created between the silencing inducing locus and the target catalase endogenes. As a positive control, the recombinant construct Y_2 was also introduced in transgenic tobacco with silenced catalase1 genes by the presence of a catalase1 antisense construct (Cat1AS in Champnongpol et al., 1996). Sixteen days after *Agrobacterium* injection, the catalase activity was determined

in protein extracts of injected leaf tissue and compared with the activity in non-injected wild type (SR1) leaf tissue (Table 2). The results indicate that domino silencing is also applicable to endogenes since the catalase activity is clearly reduced in 6 out of 7 samples, while it remains high in the negative controls. In conclusion, not only an
 5 inverted repeat-bearing silencing-inducing transgene locus, but also a silencing-inducing locus in which the two residing chimeric genes give rise to transcripts with complementarity in the 3'UTR (3'chs)(Fig.3: X₂), is able to trigger domino silencing reducing endogenous catalase expression.

10 Table 1: Results of a GUS-activity determination in protein extracts of leaf tissue harvested from tobacco plants containing different combinations of the loci X₁, Y₁ and Z₁ (Fig.2). The mean values of a number of plants (n) are given.

genotype	GUS-act. ¹ 4 weeks ² U GUS/mg TSP	N	GUS-act. Mature ² U GUS/mg TSP	n
X ₁	< ³	1	<	1
Y ₁	368 ± 165 ⁴	9	n.d.	-
Z ₁	126 ± 30	10	48 ± 8	5
X ₁ Y ₁	2 ± 1	4	4 ± 2	4
X ₁ Z ₁	139 ± 35	9	46 ± 14	5
Y ₁ Z ₁	477 ± 101	10	231 ± 106	6
X ₁ Y ₁ Z ₁ ⁵ → Y ₁ Z ₁ → X ₁ Y ₁ Z ₁	195 ± 104	16	315 ± 46	8
	4 ± 3	22	12 ± 4	9

15 ¹ The mean GUS-activity (GUS-act.) was calculated, using n samples and expressed as units (U) GUS per milligram of total soluble protein (TSP).

² The plants were analyzed in two different developmental stages; 4 weeks after sowing and at a mature stage just before onset of flowering.

³ below detection limit (1 U GUS/mg TSP)

20 ⁴ standard deviation

⁵ Growth of X₁Y₁Z₁ plants was performed in conditions that both Y₁Z₁ and X₁Y₁Z₁ plants were able to develop. A PCR screen with X₁-specific primers was performed to discriminate between presence and absence of X₁.

n.d. not determined

Table 2: Results of a catalase-activity determination in protein extracts of leaf tissue harvested from Agrobacterium injected tobacco leaves.

5

Genotype injected plant	Construct introduced via Agrobacterium injection	catalase activity 16 days after injection (60 µg TSP)
WT (SR1)	- (non-injected)	-0.2116 ² 100% ³
X ₁	PGUSchsS	-0.2556 121%
X ₁	Y ₂	-0.0589 27%
X ₁ ⁴	Y ₂	-0.0698 33%
X ₂	PGUSchsS	-0.1782 84%
X ₂	Y ₂	-0.0641 30%
X ₂	Y ₂	-0.0987 47%
X ₂ ⁴	Y ₂	-0.0914 43%
X ₂ ⁴	Y ₂	-0.1996 94%
X ₂ ⁴	Y ₂	-0.0627 30%
Cat1AS	Y ₂	-0.0439 21%

¹ X₁, see Fig. 3; X₂, see Fig. 4

² the mean of two samples independently measured (-0.2270 and -0.1963)

³ The catalase activity in wild type SR1 tobacco leaves was set to 100 %.

10 ⁴ 24 hours after Agrobacterium injection, the plants were placed under high light conditions for 24 hours (1000 µmol / m² s). This treatment is known to stimulate endogenous catalase 1 transcription. As the degree of cat suppression is similar in uninduced as in induced situation, the data indicate that enhanced transcription of the endogenous catalase target is not required to trigger domino silencing.

15

Materials and Methods

Plasmid construction

pPs35SCAT1S3chs: The T-DNA of this plasmid is schematically shown in Fig. 3 :Y₂ and the nucleotide sequence is depicted in SEQ ID N° 1.

20

Description of the transgene loci and production of hybrid plants

Locus X₁ harbours an inverted repeat about the right T-DNA border of construct pGVCHS287, carrying a neomycinphosphotransferase II (*nptII*) gene under the control of the Cauliflower mosaic virus 35S promoter (P35S) and the 3' signalling sequences of the *Anthirrinum majus* chalcone synthase gene (3'chs). The *nptII* genes are

5 posttranscriptionally silenced and can trigger in trans silencing and methylation of homologous target genes (Van Houdt et al., 2000 a and b and Fig.2).

Locus Y₁ contains a single copy of the pGUSchsS T-DNA, containing a *gus* gene under the control of P35S and 3'chs (in transformant GUSchsS29) and shows normal

10 levels of *gus* expression (Fig.2).

Locus Z₁ contains more than one copy of the pXD610 T-DNA, harbouring the *gus* gene under control of P35S and the 3'untranslated region (UTR) of the nopaline synthase gene (3'nos), (in plant LXD610-2) and shows normal *gus* expression (De Loose et al., 1995 and Fig.2).

15 Locus X₂ contains a single copy of both the pGUSchsS and pGUSchsAS T-DNA (in transformant GUSchsS+GUSchsAS 11) and triggers silencing in cis of the *gus* genes, but also in trans of (partially) homologous genes (Fig.4).

X₁ and Z₁ hemizygous plants were obtained as hybrid progeny of the crossing of tobacco plants homozygous for locus X₁ (=Holo1; Van Houdt et al., 2000 a and b) and

20 homozygous for locus Z₁ (=LXD610-2/9 De Loose et al., 1995) to wild type SR1 respectively. Y₁ hemizygous plants were obtained by crossing the hemizygous primary tobacco transformant GUSchsS29 to SR1 and selecting for the presence of locus Y₁ in the hybrid progeny. X₁Y₁ and Y₁Z₁ hemizygous plants are the hybrid progeny plants of the cross between Holo1 and GUSchsS29 and between GUSchsS29 and LXD610-2/9

25 respectively that are selected for the presence of Y₁. X₁Z₁ hemizygous plants are the hybrid progeny of the cross between Holo1 and LXD610-2/9. X₁Y₁Z₁ hemizygous plants were obtained by crossing X₁Y₁ hemizygous plants to LXD610-2/9; as we only selected for the presence of Y₁ in the hybrid progeny both Y₁Z₁ and X₁Y₁Z₁ hemizygous plants were obtained.

30

Preparation of Agrobacteria and injection

The Agrobacteria C58C1Rif^R(pGV2260)(pGUSchsS)Cb^R,PPT^R or C58C1Rif^R(pMP90)(pPs35SCAT1S3chs)Gm^R,PPT^R were mainly grown as described by Kapila et al., 1997 except that the Agrobacteria were resuspended in MMA to a final OD₆₀₀ of 1.

Greenhouse grown plants of 10 to 15 cm in height were used. Half of the third top leaf was injected via the lower surface using a 5ml syringe while the leaf remained attached to the plant. The plants were kept in the greenhouse and 16 days after injection three to four discs of 11 mm in diameter were excised from the injected tissue
5 for the preparation of a fresh protein extract to determine the catalase activity.

Enzymatic assays

Preparation of the protein extracts and GUS-activity measurements were done as previously described (Van Houdt et al., 2000 b). Preparation of the protein extracts for
10 catalase-activity measurement and the spectrophotometric catalase-activity determination was done according to Champnongpol et al., 1996.

References

- Van Houdt, H., Kovarik, A., Van Montagu, M., and Depicker, A. (2000 a). Cross-talk between posttranscriptionally silenced neomycin phosphotransferase II transgenes. *FEBS Lett.* 467, 41-46.
- Van Houdt, H., Kovarik, A., Van Montagu, M., and Depicker, A. (2000 b) Both sense and antisense RNAs are targets for the sense transgene-induced posttranscriptional silencing mechanism. *Mol. Gen. Genet.* 263, 995-1002.
- De Loose, M., Danthinne, X., Van Bockstaele, E., Van Montagu, M. and Depicker, A. , (1995) Different 5'leader sequences modulate β -glucuronidase accumulation levels in transgenic *Nicotiana tabacum* plants. *Euphytica* 85, 209-216.
- Kapila, J., De Rycke, R., Van Montagu, M. and Angenon, G. (1997) An *Agrobacterium*-mediated transient gene expression system for intact leaves. *Plant Science* 122, 101-108.
- Champhongpol, S., Willekens, H., Langebartels, C., Van Montagu, M., Inzé, D., and Van Camp, W. (1996) Transgenic tobacco with a reduced catalase activity develops necrotic lesions and induces pathogenesis-related expression under high light. *Plant J.* 10(3), 491-503.
- Thomas, C. L., Jones, L., Baulcombe, D.C. and Maule, A.J. (2001) Size constraints for targetting post-transcriptional gene silencing and for RNA-directed methylation in *Nicotiana benthamiana* using potato virus X vector. *Plant J.* 25(4), 417-425.
- De Buck, S. and Depicker, A. (2001) Disruption of their palindromic arrangement leads to selective loss of DNA methylation in inversely repeated gus transgenes in *Arabidopsis*. *Mol. Gen. Genom.* 265, 1060-1068.

25

Claims

1. A method for obtaining efficient RNA silencing of a target gene comprising the introduction of a recombinant gene into a host that comprises a silenced locus and a target gene whereby said recombinant gene comprises a region that is homologous
5 with said silenced locus and whereby said target gene has homology with said recombinant gene but has no significant homology with said silenced locus.
2. A method according to claim 1 wherein said host is a plant or plant cell.
3. A method according to claims 1 or 2 to obtain high throughput gene silencing.
4. A plant or plant cell comprising a silenced target gene obtainable by a method
10 according to claims 1 or 2.
5. A method according to claims 1 or 2 wherein said RNA silencing of the target gene is obtained in more than 95% of the hosts.
6. A method according to claims 1 or 2 wherein RNA silencing of the target gene is obtained in more than 85% of the hosts.
- 15 7. A method according to claims 1 or 2 wherein said RNA silencing of the target gene occurs at an efficiency of more than 95 % as compared to the level of the unsilenced expression of the target gene.
8. A method according to claims 1 or 2 wherein said RNA silencing of the target gene occurs at an efficiency of more than 85 % as compared to the level of the
20 unsilenced expression of the target gene.

Figure 1




Figure 2




Figure 3




Figure 4





Figure 5



SEQUENCE LISTING

<110> VLAAMS INTERUNIVERSITAIR INSTITUUT VOOR BIOTECHNOLOGIE VZW

<120> An efficient system for RNA silencing

<130> ADP/Dom/V097

<150> EP01203760.2

<151> 2001-10-05

<160> 1

<170> PatentIn version 3.1

<210> 1

<211> 10635

<212> DNA

<213> Artificial Sequence

<220>

<223> pPs35SCAT1S3chs

<400> 1
agattcgaag ctcggtcccg tgggtgttct gtcgtctcgt tgtacaacga aatccattcc 60
cattccgcgc tcaagatggc ttcccctcgg cagttcatca gggctaaatc aatctagccg 120
acttgctcgg tgaaatgggc tgcactccaa cagaaacaat caaacaacaa tacacagcga 180
cttattcaca cgcgacaaat tacaacggta tatatcctgc cagtactcgg ccgtcgaata 240
acttcgtata atgtatgcta tacgaagtta tgaattcgcg ctctatcata gatgtcgcta 300
taaacctatt cagcacaata tattgttttc attttaatat tgtacatata agtagtaggg 360
tacaatcagt aaattgaacg gagaatatta ttcataaaaa tacgatagta acgggtgata 420
tattcattag aatgaaccga aaccggcggg aaggatctga gctacacatg ctcagggttt 480
ttacaacgtg cacaacagaa ttgaaagcaa atatcatgcg atcataggcg tctcgcatat 540
ctcattaaag cagctggaag atttgatgga tcctcatcag atctcgggtga cgggcaggac 600
cggacggggc ggtaccggca ggctgaagtc cagctgccag aaaccacgt catgccagtt 660
cccgtgcttg aagccggccg cccgcagcat gccgcggggg gcatatccga gcgcctcgtg 720
catgcgcacg ctcggtcgtg tgggcagccc gatgacagcg accacgctct tgaagccctg 780
tgcctccagg gacttcagca ggtgggtgta gagcgtggag ccagtcctcg tccgctgggtg 840
gcggggggag acgtacacgg tcgactcggc cgtccagtcg taggcgttgc gtgccttcca 900

```

ggggcccgcg taggcgatgc cggcgacctc gccgtccacc tcggcgacga gccagggata 960
gcgtcccgag agacggacga ggtcgtccgt ccactcctgc ggttcctgcg gctcggtagc 1020
gaagttgacc gtgcttgtct cgatgtagtg gttgacgatg gtgcagaccg ccggcatgtc 1080
cgctcgggtg gcacggcgga tgcggccgg gcgtcgttct gggctcatgg tagatctgtt 1140
taaagcttaa cggattgaga gtgaatatga gactctaatt ggataccgag gggaatttat 1200
ggaacgtcag tggagcattt ttgacaagaa atatttgcta gctgatagtg accttaggcg 1260
acttttgaac gcgcaataat ggtttctgac gtatgtgctt agctcattaa actccagaaa 1320
cccgcggtg agtggctcct tcaatcgttg cggttctgtc agttccaaac gtaaaacggc 1380
ttgtcccgcg tcacggcggg gggtcataac gtgactccct taattctccg ctcatgatca 1440
agctacctca gcaggatccg gcgcgccatg gtcgataaga aaaggcaatt ttagatgtt 1500
aattcataac atctctcca tgacttaaaa aacttgcaaa agatttatat agaaatactt 1560
aaatattttg actaaaaaaaa aaaaaaaaaa aacacacaca taaaccaaca aataacataa 1620
attattttta tatagccttt atttcaatga tcacaacgaa acaatacaag tacaagcgt 1680
tacaagagag aaatcgccaa tatagctcac atgcagcaca catcacaata ataggaacc 1740
atgtccactt ttttattacg gaaataagaa aataacccaa ccccggtacc cgggttcata 1800
tgcttggctt cacattaagc ctagaagcta gcttttgacc cagagatttg tcagcctgag 1860
accagtatga gatccaaatg ctgcggatct cataagtgat acgaggatca gacaaggtct 1920
ccaccaccg acgaataaag cgttcttgcc tgtctggtgt gaatgagcgg tacctttctc 1980
ctggttgctt gaaattgttc tctttctgaa tgacacactt ctgcggttg ccagtcaca 2040
ttgtagaagg aataggatac ttctcagcat ggcgaacagg atcatacctt gaagggaagt 2100
agtcgatctc ctcatccctg tgcataaaat tcattggagcc atcgtagtga ttgttgtagt 2160
gagcgctttt tggagcatta gcaggtagtt gcaaatagtt tggccaagt cgatacctct 2220
gggtatcaga gtaggagaaa atacgagttt gaagcatctt atcatctgag taataaacc 2280
ctggaacaac aatagaaggg cagaaagcta gctgctcatt ctcattagag aagttatcaa 2340
tgttcttgtt cagaactaat cttccaccg gctgcaaagg caagatatcc tctggccaag 2400
ttttgtcac atcaagtga tcaaatcaa atctgtctt atgatctgga tccatagtc 2460
ccgggcagtg ggcgatttga tttaaatctc tagaatagta aattgtaatg ttgtttgtt 2520
ttgttttgtt tgtggaattt gttgtaaaaa tacggatcgt cctgcagtcc tctccaatg 2580
aaatgaactt cttatatag aggaagggtc ttgcgaagga tagtgggatt gtgcgtcatc 2640
ccttacgtca gtggagatat cacatcaatc cacttgctt gaagacgtgg ttggaacgtc 2700
ttctttttcc acgatgtccc tcgtgggtgg gggccatct ttgggaccac tgcggcaga 2760
ggcatcttga acgatagcct ttcctttatc gcaatgatgg catctttagg tgccaccttc 2820
cttttctact gtccttttga tgaagtgaca gatagctggg caatggaatc cgaggagggt 2880
tcccgatatt accctttgtt gaaaagtctc aatagccctt tggcttctg agactgtatc 2940
tttgatattc ttggagtaga cgagagtgtc gtgtccacc atgttgacga agattttctt 3000
ctgtgcatgt agtcgtaaaa gactctgtat gaactgttcg ccagtcctca cggcgagttc 3060
tgtagatcc tcgatctgaa tttttgactc catggcctt gattcagtag gaactacttt 3120
cttagagact ccaatctcta ttacttgctt tggtttatga agcaagcctt gaatcgtcca 3180

```

tactggaata gtacttctga tcttgagaaa tatatctttc tctgtgttct tgatgcagtt	3240
agtccctgaat cttttgactg catctttaac cttcttgagg aggtatttga tctcctggag	3300
attattactc gggtagatcg tcttgatgag acctgccgcg taggcctctc taaccatctg	3360
tgggtcagca ttctttctga aattgaagag gctaactctc tcattatcgg tggatgaacat	3420
ggatcgtca ctttctccgt cgaactttct tcctagatcg tagagataga gaaagtcgtc	3480
catggtgatc tccggggcaa aggagatctc tagagtcgag atttaaattcc taaatcctgc	3540
aggaagctta ccggtataac ttcgtatagc atacattata cgaagttatc catggagcca	3600
tttacaattg aatatatcct gccgccgctg ccgctttgca cccgggtggag cttgcatgtt	3660
ggtttctacg cagaactgag ccggttaggc agataatttc cattgagaac tgagccatgt	3720
gcaccttccc cccaacacgg tgagcgacgg ggcaacggag tgatccacat gggactttta	3780
aacatcatcc gtcggatggc gttgcgagag aagcagtcga tccgtgagat cagccgacgc	3840
accgggcagg cgcgcaacac gatcgcaaag tatttgaacg caggtacaat cgagccgacg	3900
ttcacggtac cggaacgacc aagcaagcta gcttagtaaa gccctcgcta gattttaatg	3960
cggatgttgc gattacttcg ccaactattg cgataacaag aaaaagccag cctttcatga	4020
tatatctccc aatttgtgta gggcttatta tgcacgctta aaaataataa aagcagactt	4080
gacctgatag tttggctgtg agcaattatg tgcttagtgc atctaacgct tgagttaagc	4140
cgcgccgga agcggcgctg gcttgaacga attgttagac attatttgcc gactaccttg	4200
tgatctcgc ctttcacgta gtggacaaat tcttccaact gatctgcgcg cgaggccaag	4260
cgatcttctt cttgtccaag ataagcctgt ctacttcaa gtatgacggg ctgatactgg	4320
gccggcaggc gctccattgc ccagtcggca gcgacatcct tcggcgcgat tttgccgggt	4380
actgcgctgt accaaatgcg ggacaacgta agcactacat ttcgctcatc gccagcccag	4440
tcgggcggcg agttccatag cgttaagggt tcatttagcg cctcaaatag atcctgttca	4500
ggaaccggat caaagagttc ctccgccgct ggacctacca aggcaacgct atgttctctt	4560
gcttttgtca gcaagatagc cagatcaatg tcgatcgtgg ctggctcgaa gatacctgca	4620
agaatgtcat tgcgctgcc ttttccaaat tgcagttcgc gcttagctgg ataagccac	4680
ggaatgatgt cgtcgtgcac aacaatgggt acttctacag cgcgagaaat ctcgctctct	4740
ccaggggaag ccgaagtttc caaaaggctg ttgatcaaag ctccgccgct tgtttcatca	4800
agccttacgg tcaccgtaac cagcaaatca atatcactgt gtggcttcag gccgccatcc	4860
actgcggagc cgtacaaatg tacggccagc aacgtcgggt cgagatggcg ctcgatgacg	4920
ccaactacct ctgatagttg agtcgatact tcggcgatca ccgcttcctt catgatgttt	4980
aactttgttt tagggcgact gccctgctgc gtaacatcgt tgctgctcca taacatcaaa	5040
catcgaccca cggcgtaacg cgcttgctgc ttggatgcc gaggcataga ctgtaccca	5100
aaaaaacagt cataacaagc catgaaacc gccactgcgc cgttaccacc gctgcgttcg	5160
gtcaagggtt tggaccagtt gcgtgagcgc atacgctact tgcattacag cttacgaacc	5220
gaacaggctt atgtccactg ggctcgtgcc ttcattccgt tccacgggtg gcgtcaccgc	5280
gcaaccttgg gcagcagcga agtcgaggca tttctgtcct ggctggcgaa cgagcgcaag	5340
gtttcgggtc ccacgcacgc tcaggcattg gcggccttgc tgttcttcta cggcaagtgc	5400

tgtgcacgga tctgcccttg cttcaggaga tcggaagacc tcggccgtcc gggcgcttgc 5460
 cgggtggtgct gaccccgat gaagtgggtc gcatcctcgg ttttctggaa ggcgagcatc 5520
 gtttgttcgc ccagcttctg tatggaacgg gcatgcggat cagtgaaggt ttgcaactgc 5580
 ggggtcaagga tctggatttc gatcacggca ccatcatcgt gcgggagggc aagggctcca 5640
 aggatcgggc cttgatgtta cccgagagct tggcaccag cctgcgcgag cagggatcga 5700
 tccaacccct ccgctgctat agtgcagtcg gcttctgacg ttcagtgcag ccgtcttctg 5760
 aaaacgacat gtcgcacaag tcctaagtta cgcgacaggc tgccgccctg cccttttctt 5820
 ggcgttttct tgcgcgtgt tttagtcgca taaagtagaa tacttgcgac tagaaccgga 5880
 gacattacgc catgaacaag agcgcgccg ctggcctgct gggctatgcc cgcgtcagca 5940
 ccgacgacca ggacttgacc aaccaacggg ccgaactgca cgcggccggc tgcaccaagc 6000
 tgttttccga gaagatcacc ggcaccaggc gcgaccgcc ggagctggcc aggatgcttg 6060
 accacctacg ccctggcgac gttgtgacag tgaccaggct agaccgcctg gcccgcagca 6120
 cccgcgacct actggacatt gccgagcgca tccaggaggc cggcgccggc ctgcgtagcc 6180
 tggcagagcc gtgggccgac accaccacgc cggccggccg catggtgttg accgtgttcg 6240
 ccggcattgc cgagttcgag cgttcctaa tcatcgaccg caccggagc gggcgcgagg 6300
 ccgccaaggc ccgaggcgtg aagtttggcc cccgcctac cctcaccgc gcacagatcg 6360
 cgcacgcccg cgagctgac gaccaggaag gccgcaccgt gaaagaggcg gctgcaactgc 6420
 ttggcgtgca tcgctcgacc ctgtaccgcg cacttgagcg cagcgaggaa gtgacgcca 6480
 ccgaggccag gcggcgcggt gccttcctg aggacgcatt gaccgaggcc gacgccctgg 6540
 cggccgccga gaatgaacgc caagaggaac aagcatgaaa ccgcaccagg acggccagga 6600
 cgaaccgttt ttcattaccg aagagatcga ggcggagatg atcgcgccg ggtacgtgtt 6660
 cgagccgccc gcgcacgtct caaccgtgcg gctgcatgaa atcctggccg gtttgtctga 6720
 tgccaagctg gcggcctggc cggccagctt ggccgctgaa gaaaccgagc gccgccgtct 6780
 aaaaagggtga tgtgtatttg agtaaaacag cttgcgtcat gcggtcgctg cgtatatgat 6840
 gcgatgagta aataa caaaa tacgcaagg gaacgcatga aggttatcgc tgtacttaac 6900
 cagaaaggcg ggtcaggcaa gacgaccatc gaacccatc tagcccgcg cctgcaactc 6960
 gccggggccg atgttctgtt agtcgattcc gatccccagg gcagtggccc cgattgggag 7020
 gccgtgcggg aagatcaacc gctaaccgtt gtcggcatcg accgcccac gattgaccgc 7080
 gacgtgaagg ccatcgccg gcgcgacttc gtagtgatcg acggagcgcc ccaggcgccg 7140
 gacttggtg tgcgcgcat caaggcagcc gacttcgtgc tgattccggt gcagccaagc 7200
 ccttacgaca tatgggccac cgcgcacctg gtggagctgg ttaagcagcg cattgaggtc 7260
 acggatggaa ggctacaagc ggcctttgtc gtgtcgccgg cgatcaaagg cacgcgcatc 7320
 ggcggtgagg ttgccgaggc gctggccggg tacgagctgc ccattcttga gtcccgtatc 7380
 acgcagcgcg tgagctaccc aggcactgcc gccgccggca caaccgttct tgaatcagaa 7440
 cccgagggcg acgtgccccg cgaggctccg gcgctggccg ctgaaattaa atcaaaactc 7500
 atttgagtta atgaggtaaa gagaaaatga gcaaaagcac aaacacgcta agtgccggcc 7560
 gtccgagcg acgcagcagc aaggctgcaa cgttgccag cctggcagac acgccagcca 7620
 tgaagcgggt caactttcag ttgccggcg aggatcacac caagctgaag atgtacgcg 7680

tacgccaagg caagaccatt accgagctgc tatctgaata catcgcgag ctaccagagt	7740
aatgagcaa atgaataaat gagtagatga attttagcgg ctaaaggagg cggcatggaa	7800
aatcaagaac aaccaggcac cgacgccgtg gaatgcccc tgtgtggagg aacgggagg	7860
tggccaggcg taagcggctg ggttgtctgc cggccctgca atggcactgg aacccccaa	7920
cccgaggaaat cggcgtgacg gtcgcaaacc atccggcccg gtacaaatcg gcgcggcgct	7980
gggtgatgac ctggtggaga agttgaaggc cgcgaggcc gccagcggc aacgcatcga	8040
ggcagaagca cggccgggtg aatcgtggca agcggccgct gatcgaatcc gcaaagaatc	8100
ccggcaaccg ccggcagccg gtgcgccgtc gattaggaag ccgccaagg gcgacgagca	8160
accagatttt ttctgtccga tgctctatga cgtgggcacc cgcgatagtc gcagcatcat	8220
ggacgtggcc gttttccgtc tgtcgaagcg tgaccgacga gctggcgagg tgatccgcta	8280
cgagcttcca gacgggcacg tagaggtttc cgcagggccg gccggcatgg ccagtgtgtg	8340
ggattacgac ctggtactga tggcgggttc ccatctaacc gaatccatga accgataccg	8400
ggaagggaag ggagacaagc ccggccgctg gttccgtcca cacgttgccg acgtactcaa	8460
gttctgcccg cgagccgatg gcggaaagca gaaagacgac ctggtagaaa cctgcattcg	8520
gttaaacacc acgcacgttg ccatgcagcg tacgaagaag gccaagaacg gccgcctgg	8580
gacggtatcc gaggggtgaag ccttgattag ccgtacaag atcgtaaaga gcgaaaccg	8640
gcggccggag tacatcgaga tcgagctagc tgattggatg taccgcgaga tcacagaagg	8700
caagaaccg gacgtgctga cggttcaccc cgattacttt ttgatcgatc ccggcatcgg	8760
ccgttttctc taccgcctgg cagcccgcg cgcaggcaag gcagaagcca gatggttgtt	8820
caagacgac tacgaacgca gtggcagcgc cggagagttc aagaagttct gtttcaccgt	8880
gcgcaagctg atcgggtcaa atgacctgcc ggagtacgat ttgaaggagg aggcggggca	8940
ggctggcccg atcctagtca tgcgtaccg caacctgac gagggcgaag catccgccgg	9000
ttcctaagt acggagcaga tgctagggca aattgcccta gcaggggaaa aaggctgaaa	9060
aggtctcttt cctgtggata gcacgtacat tgggaaccca aagccgtaca ttgggaaccg	9120
gaacccgtac attgggaacc caaagccgta cattgggaac cggtcacaca tgtaagtga	9180
tgatataaaa gagaaaaaag gcgatttttc cgctaataac tctttaaaac ttattaaaac	9240
tcttaaaacc cgctggcct gtgcataact gtctggccag cgcacagccg aagagctgca	9300
aaaagcgct acccttcggt cgctgcgct cctacgccc gccgcttcgc gtcggcctat	9360
cgcgcccgct ggccgctcaa aaatggctgg cctacggcca ggcaatctac cagggcgagg	9420
acaagccg cgtcgccac tcgaccgccc gcgcccacat caaggcacc tgcctcgcg	9480
gtttcggtga tgacggtgaa aacctctgac acatgcagct ccgggagacg gtcacagctt	9540
gtctgtaagc ggatgccggg agcagacaag cccgtcaggc cgctcagcg ggtgttgcg	9600
ggtgtcggg cgagccatg acccagtcac gtagcgatag cggagtgtat actggcttaa	9660
ctatgcggca tcagagcaga ttgtactgag agtgacacat atgcggtgtg aaataccgca	9720
cagatgcgta aggagaaaat accgcatcag gcgctcttc gcttcctcgc tcaactgactc	9780
gctgcgctc gtcgttcggc tgcggcgagc ggtatcagct cactcaaagg cggtaatag	9840
gttatccaca gaatcagggg ataacgcagg aaagaacatg tgagcaaaag gccagcaaaa	9900

ggccaggaac cgtaaaaagg ccgcgttgct ggcgtttttc cataggctcc gccccctga 9960
cgagcatcac aaaaatcgac gctcaagtca gaggtggcga aaccgacag gactataaag 10020
ataccaggcg tttccccctg gaagctccct cgtgcgctct cctgttccga ccctgccgct 10080
taccggatac ctgtccgcct ttctcccttc ggggaagcgtg gcgctttctc atagctcacg 10140
ctgtagggtat ctcagttcgg tgtaggctcg tgcctccaag ctgggctgtg tgcacgaacc 10200
ccccgttcag cccgaccgct gcgccttata cggttaactat cgtcttgagt ccaaccggt 10260
aagacacgac ttatcgccac tggcagcagc cactggtaac aggattagca gagcgaggta 10320
tgtaggcggg gctacagagt tcttgaagtg gtggcctaac tacggctaca ctagaaggac 10380
agtatttggg atctgcgctc tgctgaagcc agttaccttc ggaaaagag ttggtagctc 10440
ttgatccggc aaacaaacca ccgctggtag cggtgggtttt tttgtttgca agcagcagat 10500
tacgcgcaga aaaaaaggat ctcaagaaga tccggaaaac gcaagcgcaa agagaaagca 10560
ggtagcttgc agtgggctta catggcgata gctagactgg gcggttttat ggacagcaag 10620
cgaaccggaa ttgcc 10635

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/11188

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/82

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, PAJ, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	VAN HOUTD H ET AL: "Cross-talk between posttranscriptionally silenced neomycin phosphotransferase II transgenes" FEBS LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 467, no. 1, 4 February 2000 (2000-02-04), pages 41-46, XP004260919 the whole document	1-8
A	WO 99 53050 A (WANG MING BO ;COMMW SCIENT IND RES ORG (AU); GRAHAM MICHAEL WAYNE) 21 October 1999 (1999-10-21) cited in the application the whole document	1-8



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

13 December 2002

Date of mailing of the international search report

20/12/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kalsner, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/11188

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99 61632 A (NAP JAN PETER HENDRIK ; CPRO DLO (NL); STIEKEMA WILLEM JOHANNES (NL) 2 December 1999 (1999-12-02) cited in the application the whole document -----	1-8

INTERNATIONAL SEARCH REPORT
Information on patent family members

International Application No

PCT/EP 02/11188

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9953050	A	21-10-1999	AU 2951499 A	01-11-1999
			CA 2325344 A1	21-10-1999
			CN 1306571 T	01-08-2001
			EP 1068311 A1	17-01-2001
			WO 9953050 A1	21-10-1999
			JP 2002511258 T	16-04-2002
WO 9961632	A	02-12-1999	EP 0959133 A1	24-11-1999
			AU 4172599 A	13-12-1999
			WO 9961632 A1	02-12-1999